

T2AG rabbit pAb

Cat No.: ES9041

For research use only

Overview

Product Name T2AG rabbit pAb

Host species Rabbit WB;ELISA **Applications**

Species Cross-Reactivity Human; Mouse; Rat

Recommended dilutions WB 1:500-2000 ELISA 1:5000-20000

Immunogen Synthesized peptide derived from human protein.

at AA range: 30-110

T2AG Polyclonal Antibody detects endogenous Specificity

levels of protein.

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and

0.02% sodium azide.

Store at -20°C. Avoid repeated freeze-thaw cycles. **Storage** Transcription initiation factor IIA subunit 2 (General **Protein Name**

transcription factor IIA subunit 2) (TFIIA p12 subunit)

(TFIIA-12) (TFIIAS) (Transcription initiation factor IIA

gamma chain) (TFIIA-gamma)

Gene Name GTF2A2 TF2A2

Cellular localization Nucleus.

Purification The antibody was affinity-purified from rabbit

antiserum by affinity-chromatography using

epitope-specific immunogen.

Clonality Polyclonal Concentration 1 mg/ml **Observed band** 11kD **Human Gene ID** 2958 **Human Swiss-Prot Number** P52657

Alternative Names

Background Accurate transcription initiation on TATA-containing

> class II genes involves the ordered assembly of RNA polymerase II (POLR2A; MIM 180660) and the general initiation factors TFIIA, TFIIB (MIM 189963), TFIID (MIM 313650), TFIIE (MIM 189962), TFIIF

> (MIM 189968), TFIIG/TFIIJ, and TFIIH (MIM 189972).




+86-27-59760950 ELKbio@ELKbiotech.com



The first step involves recognition of the TATA element by the TATA-binding subunit (TBP; MIM 600075) and may be regulated by TFIIA, a factor that interacts with both TBP and a TBP-associated factor (TAF; MIM 600475) in TFIID. TFIIA has 2 subunits (43 and 12 kD) in yeast and 3 subunits in higher eukaryotes. In HeLa extracts, it consists of a 35-kD alpha subunit and a 19-kD beta subunit encoded by the N- and C-terminal regions of GTF2A1 (MIM 600520), respectively, and a 12-kD gamma subunit encoded by GTF2A2 (DeJong et al., 1995 [PubMed 7724559]).[supplied by OMIM, Mar 2008],

Western blot analysis of lysates from SW480 cells, primary antibody was diluted at 1:1000, 4° over night



+86-27-59760950

